

methylmercury in the diet. For previous reports of neuropsychological effects of methylmercury and PCP exposures see Progress in Pediatric Neurology III, PNB Publ, 1997;pp278-280.

Iatrogenic exposure to mercury after hepatitis B vaccination in preterm infants is reported by Stajich GV et al. (J Pediatr May 2000;136:679-681). Thimerosal, a mercury-derived preservative, has been used in some vaccines since the 1930s. Thimerosal is composed of 49.6% ethylmercury, which behaves like methylmercury. According to a review editorial by Pless R and Risher JF (J Pediatr May 2000;136:571-573), infants may be exposed to cumulative doses of mercury from vaccines in the first 6 months, exceeding the EPA limit of 0.1 mcg/kg/d for chronic daily exposure to methylmercury. The Advisory Committee on Immunization Practices has recommended that hepatitis B vaccine used in infants at birth should not contain thimerosal.

PRE- AND PERI-NATAL DISORDERS

EARLY PREDICTORS OF HIE ADVERSE OUTCOME

A retrospective study of 35 term infants with post-asphyxial hypoxic-ischemic encephalopathy (HIE) was conducted at the KK Women's and Children's Hospital, Singapore, to determine early predictors of mortality or major motor morbidity at 18 months of age. A severe adverse outcome occurred in 23; thirteen died and ten had major neurological sequelae. Risk factors included a low 5 min Apgar score (<4), the use of adrenaline, low arterial pH (<7.1) and high base deficit (>20 mEq/L). The high base deficit and low Apgar score combined had a positive predictive value of 100%. (Toh VC. Early predictors of adverse outcome in term infants with post-asphyxial hypoxic ischaemic encephalopathy. Acta Paediatr March 2000;89:343-347). (Respond: Dr Veronica C Toh, Department of Neonatology, KK Women's and Children's Hospital, 100 Bukit Timah Rd, Singapore).

COMMENT. The combination of a low Apgar score and high base deficit in term infants with post-asphyxial HIE is an early predictor of mortality or major neurologic sequelae.

Serum CPK and outcome of HIE. An elevated serum CPK measured within 4 hours after birth is a sensitive indicator of brain damage in asphyxiated term infants but is of limited prognostic value in assessment of neurological outcome, according to one previous report, whereas another study showed that CPK measured in cord blood correlates with outcome after asphyxia and compares favorably with imaging studies. (Progress in Pediatric Neurology I, 1991;pp 332).

Cranial ultrasonography and spectroscopy are of value in the prediction of neurodevelopmental outcome of HIE (see Progress in Pediatric Neurology II, 1994; 313-331).

MATERNAL THYROID FUNCTION AND INFANT DEVELOPMENT

The effect of maternal thyroid function in the first half of pregnancy on the neurologic development of 20 infants in the first two years of life was studied at the University of Amsterdam and Emma Children's Hospital, The Netherlands. At the age of 6 and 12 months, the mean mental developmental index (MDI) score was 16 points lower for 7 infants born to mothers with subclinical hypothyroidism compared to 6 with euthyroid mothers (P=.03 and .02, respectively). At 24 months, a mean 6 point lower MDI score was not statistically significant. One infant out of